

University of Groningen

Neuroprotective and neurodegenerative effects of the chronic expression of tumor necrosis factor α in the nigrostriatal dopaminergic circuit of adult mice

Chertoff, M.; Di Paolo, N.; Schoeneberg, A.; Depino, A.; Ferrari, C.; Wurst, W.; Pfizenmaier, K.; Eisel, U.; Pitossi, F.

Published in:
Experimental Neurology

DOI:
[10.1016/j.expneurol.2010.11.010](https://doi.org/10.1016/j.expneurol.2010.11.010)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2011

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Chertoff, M., Di Paolo, N., Schoeneberg, A., Depino, A., Ferrari, C., Wurst, W., Pfizenmaier, K., Eisel, U., & Pitossi, F. (2011). Neuroprotective and neurodegenerative effects of the chronic expression of tumor necrosis factor α in the nigrostriatal dopaminergic circuit of adult mice. *Experimental Neurology*, 227(2), 237-251. <https://doi.org/10.1016/j.expneurol.2010.11.010>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.


Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.


Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Fluoro-jade staining

30 µm coronal sections at the level of the SN were mounted onto Superfrost Plus slides and dried at room temperature for 48 hours. Slides were then immersed in a solution containing 1% NaOH in ethanol 80% (5 min), then in 70% ethanol (2 min) and finally in distilled water (2 min). Slides were transferred to a freshly prepared solution of 0.06% potassium permanganate and gently shaken for 10 min. After a rinse in distilled water for 2 min, slide-mounted sections were placed in 0.0004% Fluoro-jade B dissolved in 0.1% acetic acid in the dark for 20 min, and gently shaken. After staining, sections were washed three times in distilled water (1 min each wash), dried, immersed for 1 min in xylene and coverslipped.



Supplementary Fig. 1. Characterization of the system combining K-in mice, adenoviral vectors and the CRE/loxP system to achieve different levels of TNF- α expression in the SN. A) X-gal staining and immunohistochemistry for tyrosine hydroxylase (TH) of nigral sections injected with Adbgal. Three representative sections, at positions AP-2.8, -3.4, and -3.8 from the *bregma* according to Watson and Paxinos and encompassing the entire SN, are shown. B) Detection of nuclear CRE (left panel) and TH (center panel) and co-localization of CRE and TH (right panel) in the SN as analyzed by double immunofluorescence. C) Quantification of double-labeled TH/beta-galactosidase or TH/CRE cells throughout the entire SN relative to total TH+ cells 4 days after adenoviral transduction in that region. D) RT-PCR of the SN or hippocampus of K-in (K-in) mice or control littermates (wt) injected with adenovectors or saline in the SN. Amplification of GAPDH RNA retrotranscribed and amplified as the rest of the samples was used as a control of the amount of RNA subjected to PCR. Hip, hippocampus. Scale bar in A: 200 μ m; in B: 20 μ m.



Supplementary Fig. 2. Neurodegeneration after 100 days of AdCRE treatment. A–B) Representative SN sections stained with Fluoro-Jade-B 100 days after Adbgal (A) or AdCRE (B) injection. Arrow: degenerating neuron; arrowheads: red blood cell. Scale bar: 50 μ m.